

COPPER	GROUP	M1	FUNGICIDE
--------	-------	----	-----------



CS 2005

For Use To Control Bacteria and Fungi That Cause Spoilage On Agricultural Commodities

Ingredients

Copper Sulfate Pentahydrate*...(CAS No.7758-99-8).....	19.8%
Other Ingredients	<u>80.2%</u>
Total	100.0%

*Equivalent to 5.0% metallic copper

A Chelated Copper Product

Keep Out of Reach of Children **DANGER PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail)

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and then continue rinsing. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN or CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move a person to fresh air. If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment advice.

Mantener Fuera Del Alcance De Los Niños **PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail)

PRIMEROS AUXILLIOS	
Si entra an contacto con los ojos	<ul style="list-style-type: none">• Mantenga el ojo Abierto y enjuáguelo lenta y cuidadosamente con agua, durante 15 a 20 minutos.• Si utiliza lentes de contacto, retirelos después de los primeros 5 minutos, luego continue enjuagando [el ojo/los ojos]• Llame al centro de control de envenenamientos o a un medico para consejo de traramiento.
Si se ingiere	<ul style="list-style-type: none">• Llame inmediatamente a un centro de control de envenenamientos a un médico para consejo de tratamiento.• Si la persona puede tragar, haga que tome pequeños sorbos de agua.• No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un medico.• No administre nada por vía oral a una persona inconsciente.
Si está en la piel o en la ropa	<ul style="list-style-type: none">• Quítese la ropa contaminada• Enjuague la piel inmediatamente con bastante agua por 15-20 minutos.• Llame al centro de control de envenenamientos o a un médico para consujo de tratamiento.
Si es inhalado	<ul style="list-style-type: none">• Traslade a la persona a un lugar con aire fresco.• Si la persona no está respirando, llame al 911 o a una ambulancia inmediatamente. Luego proceda a dar respiración artificial, preferiblemente de boca a boca, si es possible.• Llame al centro de control de envenenamientos o a un medico para consejo de tratamiento.
Nota al Médico: El probable daño de las mucosas puede contraindicar el uso de un lavado gástrico.	
Cuando llame a un centro de control de envenenamientos, o a un medico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del product.	
Para emergencias médicas, / Para más información sobre al tratamiento medico de emergencia, trambien puede comunicarse, llame CHEMTREC 1-800-424-9300.	

PRECAUTIONARY STATEMENTS **Hazards to Humans and Domestic Animals**

DANGER. Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing.

DECLARACIONES de PRECAUCIÓN

Reisgos para los seres humanos y animals domésticos

PELIGRO. Corrosivo. Causa daño ocular irreversible. No permita que caiga en los ojos, la piel o la ropa.

Personal Protective Equipment

Personal Protective Equipment: Some materials that are chemical-resistant to this product are listed below. Applicators and other handlers must wear:

- Long sleeved shirt and long pants;
- Chemical resistant gloves made out of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils;
- Protective eyewear; and
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Equipo de Protección Personal (PPE, por sus siglas en inglés)

Equipo de Protección: Los aplicadores y otros manipuladores de pesticidas deben usar:

- Camisa de manga larga y pantalones largos,
- Guantes resistentes a productos químicos hechos de laminados impermeables, goma de butilo (\geq 14 mils), goma de nitrilo (\geq 14 mils), goma neoprene (\geq 14 mils), goma natural (\geq 14 mils), polietileno, cloruro de polivinilo (PVC, por sus siglas en inglés) (\geq 14 mils), o Viton® \geq 14 mils.
- Lentes o gafas protectoras
- Lentes de seguridad blindados.

Desche la ropa y otros materiales absorbentes que hayan sido empanados o muy contaminados con el concentrado de este producto. No vuelva a reutilizarlos. Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de ropa para lavar.

User Safety Requirements

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with product's concentrate. Do not re-use them.

Requerimientos de Seguridad Para el Usuario

Desche la ropa y otros materiales absorbentes que hayan sido empanados o muy contaminados con el concentrado de este producto. No vuelva a reutilizarlos. Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de ropa para lavar.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling product. As soon as possible, wash thoroughly a change into clean clothing.
- Wash the outside of gloves before removing.
- As soon as possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has potential for runoff for several months or more after application. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PELIGROS AMBIENTALES

Este pesticida es tóxico para peces e invertebrados acuáticos y puede contaminar el agua por escorrentía. Este producto puede escurrirse durante varios meses o más después de su aplicación. No vierta efluentes que contengan este producto en lagos, arroyos, estanques, estuarios, océanos ni otras aguas, a menos que cumpla con los requisitos de un permiso del Sistema Nacional de Eliminación de Descargas de Contaminantes (NPDES) y se haya notificado por escrito a la autoridad competente antes de la descarga. No vierta efluentes que contengan este producto en el alcantarillado sin notificar previamente a la autoridad local de la planta de tratamiento de aguas residuales. Para obtener orientación, comuníquese con la Junta Estatal del Agua o la Oficina Regional de la EPA

Fish Advisory Statement: This copper product is toxic to fish and aquatic organisms. Unlike most organic pesticides, copper is an element and will not break down in the environment and will therefore accumulate in sediment with repeated applications. Copper is a micronutrient, but its pesticidal application rate exceeds the amount of copper needed as a nutrient.

Declaración de advertencia sobre el consumo de pescado: Este producto de cobre es tóxico para los peces y los organismos acuáticos. A diferencia de la mayoría de los pesticidas orgánicos, el cobre es un elemento y no se descompone en el medio ambiente, por lo que se acumula en los sedimentos con aplicaciones repetidas. El cobre es un micronutriente, pero su tasa de aplicación como pesticida supera la cantidad necesaria de cobre como nutriente.

For Terrestrial Uses: This pesticide is toxic to fish and aquatic vertebrates and may contaminate water through runoff. Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

PESTICIDE RESISTANCE MANAGEMENT

For resistance management, please note that CS 2005 contains a Group M1 fungicide/bactericide. Any fungal/bacterial population may contain individuals naturally resistant to CS 2005 and other Group M1 fungicides/bactericides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of CS 2005 or other M1 fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the

manufacturer.

- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens. Report any incidence of non-performance of this product against a particular pathogen to your MAGNA-BON II, LLC retailer, representative or call 863-357-0400. If resistance is suspected, treat pathogens with a fungicide/bactericide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further reproduction.

PRODUCT INFORMATION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

INSTRUCCIONES DE USO

Es una violación de la ley federal utilizar este producto de manera inconsistente con lo indicado en su etiqueta.

USE RESTRICTIONS

- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FOR USE TO CONTROL BACTERIA AND FUNGI THAT CAUSE SPOILAGE ON AGRICULTURAL COMMODITIES: To use CS 2005 as a systemic formulation to control bacteria and fungi that cause spoilage on agricultural commodities, the following directions apply.

CS 2005 may be applied with any type of application equipment that gives uniform coverage of all foliage, including ground, aerial and low volume sprayers as specified on this label. Equipment used for application should be PVC or 316L stainless steel. CS 2005 is compatible with most fungal and insecticidal biopesticides when applied at least two (2) days before or after application of the biopesticide.

Phytotoxicity

Although CS 2005 has been tested on a wide variety of fruits, vegetables and nuts without phytotoxicity, there could be some varieties and cultivars that because of environmental factors and stages of growth could possibly foster systems. The per acre use rate of CS 2005 is applicable for dilute spraying. Depending on the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to the Minimum Recommended Spray Volume Table. Complete spray coverage is essential to assure optimum performance from CS 2005. When treating by aerial application or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

Consult CS 2005 label for specific rates and timing of application by crop. Where application rates and intervals are provided in a range (e.g., 2-4 fluid ounces and 7 to 10 days) the higher rates and shorter spray time intervals are recommended when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops. The use of a surfactant, such as Cell-U-Wett™ is acceptable for plants having waxy or hairy surfaces. CS 2005 works via surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted. CS 2005 does not produce any visible residue or have a distinct odor. It does have a residual, especially if applied with a surfactant.

Product Use Precautions

- This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
- Environmental conditions such as extended periods of wet weather, acid rain, etc., which alter the pH of the leaf surface may affect the performance of CS 2005 resulting in possible phytotoxicity or loss of effectiveness.
- It must be determined in the selection process if proper application equipment is available and if the waste associated with its use can be properly handled. Materials used on the construction of application equipment is also an important factor as agricultural chemicals are often reactive with soft metals such as aluminum and even some synthetic materials such as plastics, rubbers, etc.. Therefore it is necessary when working with equipment containing these materials, that they are thoroughly flushed with clean water after each days use.

Product Use Restrictions

- **Do not** mix with acidic compounds such as Allette™ within 14 days before or after application of same.
- **Do not** mix with pot ash.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. The PHI is 0 unless noted.

Do not enter or allow others to enter until sprays have dried.

PPE required for early entry to treated areas that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, is: coveralls, protective eyewear, chemical resistant gloves, and shoes plus socks.

ENGINEERING CONTROLS

Pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides [40 CFR 170.305.]

CONTROLES DE INGENIERÍA

Los pilotos deben utilizar una cabina cerrada que cumpla con la definición que figura en el WPS para pesticidas agrícolas [40CFR 170.305].

CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: Sprinkler including center pivot, lateral move, end row, side (wheel) roll, traveler, big gun, solid set or hand move; flood (basin); furrow; border or drip (trickle) irrigation and system(s). Do not apply this product through any other type of irrigation systems.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety device for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Posting areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds or other public facilities not including public roads, or 2) when chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive area. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters of at least 2 ½ inches tall, and all letters and the symbol shall be in a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER. This sign is in addition to any sign posted to comply with the Workers Protection Standard.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete break (air gap) between the flow outlet end of the fill pipe and the top of the overflow rim of the reservoir tank of at least

twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back toward the injection.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (i.e. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, agitation is not necessary. Adjust the pH of the water to 7 or below. If using stickers, spreaders, insecticides, nutrients, etc., add CS 2005 first. If compatibility is in question, use a compatibility jar test before mixing a whole tank. Because of a wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in the mixtures.

CS 2005 may be added through a traveling system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. CS 2005 readily disperses and needs no agitation.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, agitation is not necessary. Adjust the pH of the carrier water to 7 or below. If using stickers, spreaders, insecticides, nutrients, etc., add the CS 2005 first. If compatibility is in question, use a compatibility jar test before mixing a whole tank. Because of a wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in the mixtures.

CS 2005 may be added through a traveling irrigation system or at the last 30 minutes of solid set or hand moved irrigation systems. CS 2005 readily disperses and needs no agitation.

FLOOR (BASIN), FURROW AND BORDER CHEMIGATION

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back flow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination back flow.
- b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back toward the injection pump.
- c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- e. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- f. Systems must use a metering pump, such as a positive displacement pump (i.e. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

When mixing, agitation is not necessary. Adjust the pH of the carrier water to 7 or below. If using stickers, spreaders, insecticides, nutrients, etc., add the CS 2005 first. If compatibility is in question, use a compatibility jar test before mixing a whole tank. Because of a wide variety of possible combinations which can be encountered, observe all cautions and limitations on the labels of all products used on the mixtures.

CS 2005 may be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. CS 2005 readily disperses and needs no agitation.

DRIP (TRICKLE) CHEMIGATION

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of liquid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor pump stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump

motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (i.e. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

When mixing, agitation is not necessary. Adjust the pH of the carrier water to 7 or below. If using stickers, spreaders, insecticides, nutrients, etc., add the CS 2005 first. If compatibility is in question, use a compatibility jar test before mixing a whole tank. Because of a wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in the mixtures. CS 2005 may be added through a traveling irrigation system continuously or at the last 30 of solid set or hand moved irrigation systems. CS 2005 readily disperses and needs no agitation.

TANK MIXING

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involving tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture

FOR SPRAY AND SOIL DRENCH APPLICATIONS

Always spray for total foliage coverage. When re-spraying the rates and severity of the disease vary with unforeseen conditions. However, in the event of severe disease, spraying intervals can be shortened to 3 to 5 days. At times, lower rates can be as effective as higher rates and should be tried first. Usually, preventive programs may be maintained at lower rates. Use of low volume spraying is effective against Botrytis and not effective against established powdery mildew and Xanthomonas infections. Also, applications on actively growing tissue may be more effective than applications on dormant tissue.

MINIMUM RECOMMENDED SPRAY VOLUME (GALLONS) PER ACRE WHEN APPLYING CS 2005

GROUND			
CROP	AERIAL	DILUTE	CONCENTRATE*
Vegetables	3	20	30
Field Crops	3	20	30
Small Fruits	5	150	30
Vines	5	150	30
Tree Crops	10	400	50
Citrus	10	125	30

*Pesticide application equipment such as Curtec® or other similar sprayers which are capable of obtaining coverage at low volumes may be used at as low as 20 gpa of spray volume.

The following specific instructions are based on general application procedures. The recommendations of the State Extension Service should be closely followed as to timing, frequency and numbers of sprays per season.

SPRAY DRIFT

For Aerial Application:

1. Do not release spray at a height greater than 10 ft above the vegetative canopy or water, unless a greater application height is necessary for pilot safety.
2. Applicators are required to use a medium or coarse droplet size (ASABE S572.1).
3. Do not apply when wind speed exceeds 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
4. Applicators must use ½ swath displacement upwind at the downwind edge of the application area.
5. Do not apply during temperature inversions.

For Ground Boom Application:

1. Apply with the spray release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
2. Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
3. Do not apply when wind speeds exceed 15 miles per hour at the application site.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance Of Droplet Size:

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume: Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure: Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle: Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust Nozzles: Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

Boom Height – Ground Boom:

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height – Aircraft:

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

Shielded Sprayers:

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature And Humidity:

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions:

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind:

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

**FROST INJURY PROTECTION
BACTERIAL ICE NUCLEATION INHIBITOR**

Application of CS 2005 made to all crops listed on this label at rates and stages of growth indicated on this label, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas flourescens*) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

The chart below is used to calculate correct ppm* of active ingredient (A.I.) per volume of carrier water. These volumes can be used for either conventional or concentrated sprays.

**AMOUNT OF CS 2005 PER VOLUME OF WATER
FOR PROPER PPM* OF APPLIED ACTIVE INGREDIENT (AS COPPER)**

ppm* of A.I.	30 gallons per acre	50 gallons per Acre	100 gallons per Acre	125 gallons per Acre	250 gallons per Acre	500 gallons per Acre
50	3.85 oz.	6.4 oz.	12.8 oz.	16 oz.	32 oz.	64 oz.
75	5.78 oz.	9.6 oz.	19.2 oz.	24 oz.	48 oz.	96 oz.
100	7.70 oz.	12.8 oz.	25.6 oz.	32 oz.	64 oz.	1 gal.
125	9.6 oz.	16 oz.	32 oz.	40 oz.	80 oz.	1.25 gal.
150	11.5 oz.	19.2 oz.	38.4 oz.	48 oz.	96 oz.	1.5 gal.
200	15.4 oz.	25.6 oz.	51.2 oz.	64 oz.	1 gal.	2 gal.
250	19.2 oz.	32 oz.	64 oz.	80 oz.	1.25 gal.	2.5 gal.

* ppm = parts per million (as copper)

CITRUS

Grapefruit, Kumquat, Lemon, Lime, Orange, Tangelo, Tangerine

Disease	Rate per Acre per 100 gal. of water	ppm copper per 100 gallons of water	Instructions
Brown Rot	30-70 oz.	120-275 ppm	Apply at first indication of rain or first appearance of Brown Rot. Re-apply as needed during wet weather.
Greasy Spot, Pink Pitting	25.6-64 oz.	100-250 ppm	Apply during mid-summer.
Scab	25.6-64 oz.	100-250 ppm	Apply shortly before trees begin to flush. Re-apply at 2/3 petal fall. Re-apply 4 weeks later, if necessary.
Melanose	25.6-64 oz.	100-250 ppm	Apply 2 times per year before the onset of spring and autumn rains.
Canker (Suppression)	12.8-64 oz.	50-250 ppm	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require additional applications. Number and timing of applications will be dependent on disease pressure. Under heavy pressure, each new flush of growth should be sprayed. Heavily infected trees should be sprayed with a minimum dosage of 250 ppm with a follow up spray after 7-14 days at 200 ppm.
<p>Do not apply more than 3.15 lbs Cu²⁺ per acre per application.</p> <p>Do not apply more than 12.6 lbs Cu²⁺ per acre per year.</p> <p>Do not make applications less than 7 days apart.</p>			

FIELD CROPS

Crop	Disease	Rate/Acre per 100 gallons of water	ppm copper per 100 gallons of water	Instructions
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	19.2-32 oz.	7.5-125 ppm	Apply 10 to 14 days before each harvest or earlier if disease threatens. IMPORTANT: Spray injury may occur with sensitive varieties such as Lahontan. Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 1.12 lb. Cu ²⁺ per acre per year. Do not make applications less than 30 days apart.
Corn (Field Corn, Popcorn, Sweet corn)	Bacterial Stalk Rot	19.2-32 oz.	75-125 ppm	Begin treatment when disease first appears and repeat every 7 to 10 days or as needed. Use the higher rates and shorter spray intervals when conditions favor disease. Do not apply more than 1.05 lb. Cu ²⁺ per acre per application. Do not apply more than 4.2 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.
Peanut	Cercospora leaf	19.2-25.6 oz.	75-100 ppm	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 10 to 14 day intervals or as needed. Reduce sprays to
				7 day intervals during humid weather. Use the higher rates when conditions favor disease. Do not apply more than 0.79 lb. Cu ²⁺ per acre per application. Do not apply more than 4.74 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.

Potato	Early Blight, Late	19.2-32 oz.	75-125 ppm	Apply 75 to 125 ppm at 7 to 10 day intervals or as needed starting when plants are 2 to 6 inches high in locations where disease is light. Add up to 32 oz. per acre when disease is more severe.
	Do not apply more than 2.5 lb. Cu ²⁺ per acre per application. Do not apply more than 25 lb. Cu ²⁺ per acre per year. Do not make applications less than 5 days apart.			
Sugar Beets	Cercospora Leaf	19.2-38.4 oz.	75-150 ppm	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals or as needed. Use higher rates when conditions favor disease. Addition of a sticker/spreader is recommended.
	Do not apply more than 1.31 lb. Cu ²⁺ per acre per application. Do not apply more than 7.86 lb. Cu ²⁺ per acre per year. Do not make applications less than 10 days apart.			
Wheat, Barley, Oats	Helminthosporium Spot Blotch, Septoria Leaf Blotch	19.2-25.6 oz.	75-100 ppm	Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease.
	Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 1.06 lb. Cu ²⁺ per acre per year. Do not make applications less than 10 days apart.			

SMALL FRUITS

Crop	Disease	Rate/Acre per 100 gallons, of water	ppm copper per 100 gallons of water	Instructions
Blackberry (Aurora, Boysen, Cascade, Chehalem, Logan, Marion, Santiam, Thornless	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	High 32 oz.	125 ppm	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	Low 19.2 oz.	75 ppm	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural- type spray oil may be added.
Evergreen)			IMPORTANT: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.	
	Do not apply more than 2.9 lb. Cu ²⁺ per acre per application. Do not apply more than 10.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			

Blueberry	Bacterial Canker	32-51.2 oz.	125-200 ppm	Make application before fall rains and a second application 4 weeks later. Use the higher rates when conditions favor disease.*
	Fruit Rot, Phomopsis Twig Blight	25.6-51.2 oz.	100-200 ppm	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 10 to 14 day intervals or as needed before blooms open.
	Do not apply more than 2.1 lb. Cu ²⁺ per acre per application. Do not apply more than 8.4 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Cranberry*	Fruit Rot	51.2 oz.	200 ppm	Make application in late bloom. Apply one or two additional applications at 10 to 14 day intervals or as needed depending on disease severity.
	Rose Bloom	51.2 oz.	200 ppm	Apply three sprays on 10 to 14 day schedule or as needed as soon as symptoms are observed.
	Bacterial Stem Canker	51.2 oz.	200 ppm	Apply to vines after harvest and again in spring at bud swell. Apply one or two additional applications at 10 to 14 intervals or as needed depending on disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (<i>Monilinia</i>)	51.2 oz.	200 ppm	Apply delayed dormant spray in the spring. Repeat at 10 to 14 day intervals or as needed through pre-bloom.
	Do not apply more than 2.1 lb. Cu ²⁺ per acre per application. Do not apply more than 6.3 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)			
Currant*, Gooseberry	Anthracnose, Leaf Spot	64 oz.	250 ppm	Make Initial application after first leaves have expanded. Continue on a 10 to 14 day schedule or as needed during wet conditions in the spring. Make an additional application after harvest.
	Do not apply more than 10.0 lb. Cu ²⁺ per acre per application. Do not apply more than 2.5 lb. Cu ²⁺ per acre per year. Do not make applications less than 10 days apart. (*Not Permitted in California)			
Raspberry	Anthracnose, Cane Spot, Leaf Spot, <i>Pseudomonas</i> Blight, Purple Blotch, Yellow Rust	High 32 oz.	125 ppm	Make fall application after harvest. Apply delayed dormant spray after training in the spring. If needed, agricultural-type spray oil may be added.

	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	Low 19.2 oz.	75 ppm	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural-type spray oil may be added. IMPORTANT: Crop injury may occur if applied to foliage under certain environmental conditions
--	---	--------------	--------	--

				such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.
				Do not apply more than 2.0 lb. Cu ²⁺ per acre per application. Do not apply more than 10.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.
Strawberry	Angular Leaf Spot (<i>Xanthomonas</i>), Leaf Blight, Leaf Scorch, Leaf Spot	19.2-25.6 oz.	75-100 ppm	Begin application when plants are established and continue on a weekly schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease. IMPORTANT: Discontinue applications if signs of crop injury appear.
				Do not apply more than 1.0 lb. Cu ²⁺ per acre per application or 1.5 lb. Cu ²⁺ per acre per application for severe disease. Do not apply more than 6.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.

TREE CROPS

Crop	Disease	Rate/Acre per 100 gallons of water	ppm copper per 100 gallons of water	Instructions
Almond, Apricot, Cherry, Plum Prune	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	32-64 oz.	200-250 ppm	Make first application before fall rains and a second at late dormant. Use the higher rates when conditions favor disease. If needed, agricultural-type spray oil may be added. For Cherries: Where disease is severe, an additional application shortly after harvest may be required. IMPORTANT: Foliar injury may occur from post-bloom sprays on almonds, especially on NePlus varieties.

	Blossom Brown Rot, Coryneum Blight (Shot Hole)	Almond: 51.2-64 oz. All others: 60-90 oz.	200-250 ppm	Apply during early bloom. Do not apply after full bloom or injury may occur. Use the higher rates when rainfall is heavy and disease pressure is high.
--	--	--	-------------	--

	Black Knot* (Plum)	32-64 oz.	125-250 ppm	Make application at bud swell up to early bloom for early disease suppression. Apply before full bloom. Use the higher rates when rainfall is heavy and disease pressure is high. IMPORTANT: To avoid plant
--	--------------------	-----------	-------------	--

	Cherry Leaf Spot* (Sour Cherries Only)	38.4-64 oz.	150-250 ppm	injury, do not use after full bloom. Apply at petal fall as well as 1 to 2 times after petal fall. Use the lower rates where disease infection is light and use the higher rates for a dormant application or where disease infection is moderate to heavy. Do not apply to sweet cherry or the English Morello variety as severe injury will result. IMPORTANT: Moderate to severe injury such as leaf spotting and defoliation may occur from post bloom applications.
Do not apply more than 1.5 lb. Cu ²⁺ per acre per application during the bloom/growing season. Do not apply more than 8.0 lb. Cu ²⁺ per acre per application during the dormant/late dormant season. Do not apply more than 18.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart during the dormant/late dormant season and no less than 5 days apart during the bloom/growing season.				
Apple	Anthracnose, Blossom Blast, European Canker (<i>Necria</i>), Shoot Blast (<i>Pseudomonas</i>)	51.2-64 oz.	200-250 ppm	Apply before fall rains. Use the higher rates when conditions favor disease. IMPORTANT: Use on yellow varieties may cause discoloration. To avoid discoloration, pick before spraying.

	Apple Scab, Fire Blight	51.2-64 oz.	200-250 ppm	<p>Make application between silver-tip and green-tip. Apply as a full cover spray for early season disease suppression.</p> <p>Extend spray schedule where fruit finish is not a concern: Continued applications may be made at 5 to 7 day intervals or as needed.</p> <p>IMPORTANT: Moderate to severe crop injury may occur from late application.</p> <p>Moderate to severe crop injury may result from an extended spray schedule. It is not intended for fresh market apples or for apples where fruit finish is a concern as it is likely to cause fruit russetting.</p>
	Collar Rot, Crown Rot	32 oz.	125 ppm	Mix 100 gallons of water. Apply 4 gallons of solution as a drench

				on the lower trunk area of each tree. Apply in early spring or in fall after harvest for best results. Do not apply to foliage or fruit.
				<p>Do not apply more than 0.5 lb. Cu²⁺ per acre per application during the bloom/growing season.</p> <p>Do not apply more than 6.0 lb. Cu²⁺ per acre per application during the Fall/ dormant season.</p> <p>Do not apply more than 16.0 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 5 days apart during the bloom/growing season.</p> <p>Only one application is permitted during the Fall/ dormant season.</p>
Avocado	Anthracnose, Blotch, Scab	51.2-64 oz.	200-250 ppm	Apply when bloom buds begin to swell and continue application at monthly intervals for five to six applications. Use the higher rates when conditions favor disease.
				<p>Do not apply more than 3.15 lb. Cu²⁺ per acre per application.</p> <p>Do not apply more than 18.9 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 14 days apart.</p>
Banana*	Sigatoka (Black and Yellow)	19.2 oz.	75 ppm	<p>Apply by air in 3 gallons of water. If needed, agricultural-type spray oil may be added. Apply on a 14 day schedule or as needed throughout the wet season. Apply at 21 day intervals or as needed during dry periods.</p>
	Black Pitting	32 oz.	125 ppm	Mix 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.

	Do not apply more than 1.05 lb. Cu ²⁺ per acre per application. Do not apply more than 18.9 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)			
Cacao*	Black Pod	19.2-64 oz.	75-250 ppm	Begin applications at the start of the rainy season and continue while infection conditions persist.
	Do not apply more than 3.15 lb. Cu ²⁺ per acre per application. Do not apply more than 15.75 lb. Cu ²⁺ per acre per year. Do not make applications less than 14 days apart. (*Not Permitted in California)			
Coffee	Coffee Berry Disease (<i>Collectotrichum coffeatum</i>)	38.4-64 oz.	150-250 ppm	Apply first spray after flowering and before onset of long rains and then at 21 to 28 day intervals or as needed until picking. Use the higher rates when conditions favor disease.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	38.4-64 oz.	150-250 ppm	Begin spray program before the onset of long rainy periods and continue throughout the rainy season at 14 to 21 day intervals or as needed. The critical time for spraying to control disease is just before, during and after flowering(s), especially when coinciding with wet weather. Use the higher rates when rainfall is

				heavy and disease pressure is high.
	Leaf Rust (<i>Hemileia</i>)	19.2-32 oz.	75 ppm	Apply before the onset of rain and then at 21 day intervals or as needed while the rains continue. Use the higher rates when rainfall is heavy and disease pressure is high.
	Iron Spot. (<i>Cercospora coffeicola</i>), Pink Disease (<i>Corticium salmonicolor</i>)	19.2 oz.	75 ppm	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
	Do not apply more than 2.1 lb. Cu ²⁺ per acre per application. Do not apply more than 12.6 lb. Cu ²⁺ per acre per year. Do not make applications less than 14 days apart.			
Filbert*	Bacterial Blight	64-115.2 oz.	250-450 ppm	Apply as a spray to trees after harvest. In seasons of heavy rainfall, apply a second spray when three fourths of the leaves have dropped. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil may be added. (Permitted only in Washington State and Oregon)

	Eastern Filbert Blight	64-115.2 oz.	250-450 ppm	Apply as a dilute spray in adequate water for thorough coverage. Make applications starting at bud swell to bud break and continue at 2 week intervals or as needed until early May. Thorough coverage is essential. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil may be added.
Do not apply more than 6.0 lb. Cu ²⁺ per acre per application. Do not apply more than 18.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 14 days apart. (*Not Permitted in California)				
Mango*	Anthracnose	38.4-64 oz.	150-250 ppm	Apply monthly after fruit set until harvest. Use the higher rates when rainfall is heavy and disease pressure is high.
Do not apply more than 2.6 lb. Cu ²⁺ per acre per application. Do not apply more than 18.2 lb. Cu ²⁺ per acre per year. Do not make applications less than 30 days apart. (*Not Permitted in California)				
Olive	Olive Knot, Peacock Spot	64-76.8 oz.	250-300 ppm	Make first application before winter rains begin. A second application in early spring should be made if disease is severe. Apply the higher rates for heavy
disease pressure or when conditions favor disease development.				
Do not apply more than 3.15 lb. Cu ²⁺ per acre per application. Do not apply more than 6.3 lb. Cu ²⁺ per acre per year. Do not make applications less than 30 days apart..				
Peach, Nectarine	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	51.2-76.8 oz..	200-300 ppm	Make first application before fall rains and a second at late dormant. For peach leaf curl, late dormant application must be made before leaf buds swell. Use the higher rates when rainfall and disease pressure is high. If needed, agricultural-type spray oil may be added.
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	51.2-76.8 oz.	200-300 ppm	Full cover spray at pink bud. Use the higher rates when conditions favor disease.
	Bacterial Spot	19.2 oz.	75 ppm	Post-bloom application applied at first and second cover sprays. RESTRICTION: do not spray 3 weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays.

	<p>Do not apply more than 8.0 lb. Cu²⁺ per acre per application during the dormant/late dormant season.</p> <p>Do not apply more than 1.5 lb. Cu²⁺ per acre per application during the bloom/ growing season.</p> <p>Do not apply more than 18.0 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 7 days apart during the dormant/late dormant season and no less than 5 days apart during the bloom/ growing season.</p> <p>(*Not Permitted in California)</p>			
Pear*	Fire Blight	19.2 oz.	75 ppm	Apply 5 day intervals or as needed throughout the bloom period. IMPORTANT: Russetting may occur in copper sensitive varieties. Excessive dosages may cause fruit russet on any variety.
	Blossom Blast (<i>Pseudomonas</i>)	51.2-76.8 oz.	200-300 ppm	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rates when disease pressure is high or when conditions favor disease development.
	<p>Do not apply more than 0.8 lb. Cu²⁺ per acre per application during the Fall.</p> <p>Do not apply more than 0.5 lb. Cu²⁺ per acre per application during the bloom/ growing season.</p> <p>Do not apply more than 16.0 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 5 days apart during the bloom/growing season.</p> <p>Only 1 application per season may be made in the Fall.</p> <p>(*Not Permitted in California)</p>			
Pecan*	Kernel Rot, Shuck Rot (<i>Phytophthora cactorum</i>), Zonate Leaf Spot	19.2-32 oz.	75-125 ppm	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to

	(<i>Cristulariella pyramidalis</i>)			4 week intervals or as needed, starting at kernel growth and continue until shucks open. Use the higher rates and shorter spray intervals if frequent rainfall occurs.
	Ball Moss, Spanish Moss	38.4-64 oz.	150-250 ppm	Apply in 100 gallons of water in the spring when ball moss is actively growing, using 1 ½ gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.
<p>Do not apply more than 2.1 lb. Cu²⁺ per acre per application.</p> <p>Do not apply more than 6.3 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 14 days apart.</p> <p>(*Not Permitted in California)</p>				
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria alternata</i>), Septoria Leaf Blight	32-64 oz.	125-250 ppm	Make initial application at bud swell and repeat on a 14 to 28 day schedule or as needed. If disease conditions are severe, use the higher rates and shorter spray intervals.

	<p>Do not apply more than 2.1 lb. Cu²⁺ per acre per application.</p> <p>Do not apply more than 8.4 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 14 days apart.</p>			
Quince*	Fire Blight	19.2 oz.	75 ppm	Apply at 5 day intervals or as needed throughout the bloom period. Apply in adequate water for thorough coverage. (Permitted only in Washington State and Oregon)
	<p>Do not apply more than 0.5 lb. Cu²⁺ per acre per application during the bloom/ growing season.</p> <p>Do not apply more than 8.0 lb. Cu²⁺ per acre per application during the Fall.</p> <p>Do not apply more than 16.0 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 5 days apart during the bloom/growing season.</p> <p>Only 1 application per season may be made in the Fall.</p> <p>(*Not Permitted in California)</p>			
Walnut	Walnut Blight	32-51 oz..	125-200 ppm	<p>Apply at first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage or as needed when frequent rainfall or extended periods of moisture occur. Thorough coverage of catkins, leaves and nutlets is essential for effective control.</p> <p>IMPORTANT: Adequate control may not be obtained when copper tolerant species of <i>Xanthomonas</i> bacteria are</p>

			present.
	<p>Do not apply more than 3.15 lb. Cu²⁺ per acre per application.</p> <p>Do not apply more than 25.2 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 7 days apart.</p>		

VEGETABLES

Crop	Disease	Rate/Acre per 100 gallons of water	ppm copper per 100 gallons of water	Instructions
Bean (Dry, Green)*	Brown Spot, Common Blight, Halo Blight	19.2-25.6 oz.	75-100 ppm	For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule or as needed depending on environmental conditions. Use the higher rates for more severe disease.
	<p>Do not apply more than 2.0 lb. Cu²⁺ per acre per application.</p> <p>Do not apply more than 4.74 lb. Cu²⁺ per acre per year.</p> <p>Do not make applications less than 7 days apart.</p> <p>(*Not Permitted in California)</p>			

Beet (Table Beet, Beet Greens)*	Cercospora Leaf Spot	19.2-32 oz.	75-125 ppm	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals or as needed. Use the higher rates when conditions favor disease.
	Do not apply more than 1.31 lb. Cu ²⁺ per acre per application. Do not apply more than 7.86 lb. Cu ²⁺ per acre per year. Do not make applications less than 10 days apart. (*Not Permitted in California)			
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	19.2 oz.	75 ppm	Begin applications when disease first threatens and repeat at 7 to 14 day intervals or as needed depending on disease severity.
	Do not apply more than 1.0 lb. Cu ²⁺ per acre per application. Do not apply more than 5.0 lb. Cu ²⁺ per acre per year. The minimum retreatment interval is 7 days.			
Celery, Celeriac	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	19.2 oz.	75 ppm	Begin applications as soon as plants are first established in the field, repeating at 5 to 7 day intervals or as needed depending on disease severity and environmental conditions.
	Do not apply more than 1.0 lb. Cu ²⁺ per acre per application. Do not apply more than 5.3 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Crucifers (Broccoli, Brussels Sprout, Cabbage, Cauliflower, Collard Greens,	Black Leaf Spot (<i>Alternaria</i>), Black Rot (<i>Xanthomonas</i>), Downy Mildew	19.2-25.6 oz.	75-100 ppm	Begin application after transplants are set in the field, or shortly after emergence of field seeded crops or when conditions favor disease development. Apply at 7 to 10 day intervals or as needed. Use the higher rates when conditions favor disease.
Mustard Greens, Turnip Greens)				IMPORTANT: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.
	Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 2.65 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon)	Alternia Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (suppression)	19.2-25.6 oz.	75-100 ppm	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat at 5 to 7 day intervals or as needed. Use the higher rates when conditions favor disease. IMPORTANT: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs.
	Do not apply more than 1.05 lb. Cu ²⁺ per acre per application. Do not apply more than 5.25 lb. Cu ²⁺ per acre per year. Do not make applications less than 5 days apart.			

Eggplant	Alternaria Blight, Anthracnose, Phomopsis	19.2 oz.	75 ppm	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals or as needed depending on disease severity.
	Do not apply more than 0.79 lb. Cu ²⁺ per acre per application. Do not apply more than 7.9 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Okra*	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	19.2-32 oz.	75 ppm	Begin treatment when disease first threatens and repeat every 5 to 10 days or as needed depending on disease severity. Use the higher rates and shorter spray intervals when conditions favor disease.
	Do not apply more than 1.05 lb. Cu ²⁺ per acre per application. Do not apply more than 5.25 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)			
Onion, Garlic	Bacterial Blight, Downy Mildew, Purple Blotch	19.2 oz.	75 ppm	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals or as needed depending on disease severity. Can cause phytotoxicity to leaves.
	Do not apply more than 1.0 lb. Cu ²⁺ per acre per application. Do not apply more than 6.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Pea	Powdery Mildew	19.2-25.6 oz.	75-100 ppm	Begin applications when disease symptoms first appear and repeat weekly intervals or as needed. Use the higher rates when conditions favor disease.
	Do not apply more than 0.79 lb. Cu ²⁺ per acre per application. Do not apply more than 3.95 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			
Pepper	Anthracnose, Bacterial	19.2- 25.6 oz.	75-100 ppm	Begin applications when
	Spot, Cercospora Leaf Spot			
	conditions first favor disease development and repeat at 7 to 10 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease.			
	Do not apply more than 0.79 lb. Cu ²⁺ per acre per application. Do not apply more than 11.85 lb. Cu ²⁺ per acre per year. Do not make applications less than 9 days apart.			
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, White Rust	19-25.6 oz.	75-100 ppm	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals or as needed. Use the higher rates when conditions favor disease. IMPORTANT: Flecking may occur on spinach leaves.
	Do not apply more than 0.79 lb. Cu ²⁺ per acre per application. Do not apply more than 3.95 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			

Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	19.2-32 oz.	75-125 ppm	Begin applications when disease first threatens and repeat at 5 to 10 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease.
	Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 17.4 lb. Cu ²⁺ per acre per year. Do not make applications less than 3 days apart.			
Watercress	Cercospora, Leaf Spot	19.2 oz.	75 ppm	For applications made to watercress, production fields must be drained of water at least 24 hours prior to each application and water must not be reapplied to the field for a minimum of 24 hours following each application. Copper must not be applied to watercress during the aquatic production phase. Begin applications when plants are first established in the field, repeating at 7 to 14 day intervals or as needed depending on disease severity. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.
	Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 2.12 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart.			

VINES

Crop	Disease	Rate/Acre per 100 gallons of water	ppm copper per 100 gallons of water	Instructions
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	19.2-32 oz.	75-125 ppm	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. Use the higher rates when conditions favor disease. IMPORTANT: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosette. For table grapes, spraying up until harvest is acceptable, for all others discontinue spray 15 days before harvest.

	Do not apply more than 3.0 lb. Cu ²⁺ per acre per application. Do not apply more than 20.0 lb. Cu ²⁺ per acre per year. Do not make applications less than 3 days apart.			
Hops*	Downy Mildew	19.2 oz.	75 ppm	Make crown treatments after pruning, but before training. After Training, additional treatments are needed at about 10 day intervals. RESTRICTION: Discontinue use two weeks before harvest.
	Do not apply more than 0.53 lb. Cu ²⁺ per acre per application. Do not apply more than 2.65 lb. Cu ²⁺ per acre per year. Do not make applications less than 10 days apart. (*Not Permitted in California)			
Kiwi*	<i>Erwinia herbicola</i> , <i>Pseudomonas flourescens</i> , <i>Pseudomonas syringae</i>	38.4 oz.	150 ppm	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of three applications may be made.

MISCELLANEOUS

Crop	Disease	Rate/Acre per 100 gallons of water	ppm copper per 100 gallons of water	Instructions
Atemoya*	Anthracnose	25.6-38.4 oz.	100-150 ppm	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher
	rates for severe disease. Do not apply more than 3.15 lb. Cu ²⁺ per acre per application. Do not apply more than 12.6 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)			
Carambola*	Anthracnose	38.4-51.2 oz.	150-200 ppm	Make initial application before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.
	Do not apply more than 2.1 lb. Cu ²⁺ per acre per application. Do not apply more than 10.5 lb. Cu ²⁺ per acre per year. The minimum retreatment interval is 7 days. (*Not Permitted in California)			
Chives	Downy Mildew	19.2 oz.	75 ppm	Begin application when plants are established in the field. Repeat every 7 to 10 days or as needed depending on disease conditions.

	<p>Do not apply more than 0.53 lb. Cu²⁺ per acre per application. Do not apply more than 2.65 lb. Cu²⁺ per acre per year. Do not make applications less than 7 days apart.</p>			
Dill	Phoma Leaf Spot, Rhizoctonia Foliage Blight	19.2-25.6 oz	75-100 ppm	Begin applications when plants are first established in the field and repeat at 7-10 day intervals or as needed depending on disease severity and environmental conditions. Use the higher rates for severe disease.
	<p>Do not apply more than 0.79 lb. Cu²⁺ per acre per application. Do not apply more than 3.95 lb. Cu²⁺ per acre per year. Do not make applications less than 7 days apart.</p>			
Guava*	Anthracnose, Red Algae	25.6-38.4 oz.	100-150 ppm	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.
	<p>Do not apply more than 1.23 lb. Cu²⁺ per acre per application. Do not apply more than 4.92 lb. Cu²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)</p>			
Litchi*	Anthracnose	25.6-38.4 oz.	100-150 ppm	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Use the higher rates for severe disease.
	<p>Do not apply more than 1.23 lb. Cu²⁺ per acre per application. Do not apply more than 4.92 lb. Cu²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)</p>			
Macadamia*	Anthracnose	38.4-64 oz.	150-250 ppm	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher

				rates for severe disease.
	Phytophthora Blight (<i>P. capsici</i>), Raceme Blight (<i>Botrytis cinerea</i>)	38.4-64 oz.	150-250 ppm	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage. Use the higher rates when conditions favor disease.
	<p>Do not apply more than 2.36 lb. Cu²⁺ per acre per application. Do not apply more than 9.44 lb. Cu²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)</p>			
Mamey Sapote*	Algal Leaf Spot	38.4-64 oz.	150-250 ppm	Apply when conditions favor disease development. Repeat on a 14 to 30 schedule or as needed as disease severity and environmental conditions dictate. Use the higher rates when conditions favor disease.

	Do not apply more than 2.1 lb. Cu ²⁺ per acre per application. Do not apply more than 8.4 lb. Cu ²⁺ per acre per year. Do not make applications less than 14 days apart. (*Not Permitted in California)			
Pitahaya (Dragon fruit)*	<i>Xanthomonas campestris, Dothiorella</i>	32-51 oz.	125-200 ppm	Begin applications at the start of the rainy season and continue while infection conditions persist.
				Do not apply more than 1.23 lb. Cu ²⁺ per acre per application. Do not apply more than 4.92 lb. Cu ²⁺ per acre per year. Do not make applications less than 7 days apart. (*Not Permitted in California)

GREENHOUSE and SHADEHOUSE CROPS

Notice to Users: CS 2005 may be used in greenhouses and shade houses to control diseases on crops which appear on this label, and specific instructions have been developed for crops listed. The grower should bear in mind the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions. Neither the manufacturer nor the seller has determined whether or not CS 2005 can be used safely on all green house and shadehouse grown crops. Consequently, injury arising from the use of CS 2005 on these types of greenhouse and shadehouse crops is the responsibility of the user. The user should determine whether CS 2005 can be used safely prior to commercial use. In a small area, apply the specified rates to the plants in question, i.e. foliage, fruit etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

Apply 100 gallons of CS 2005 per acre. Apply CS 2005 according to specific rates given for those crops in ounces (mL) per 100 gallons per acre. CS 2005 should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat at 7 to 14 day intervals or as needed; use shorter spray intervals during periods when severe disease conditions persist.

NOTE: Phytotoxicity may occur on young tender flush when CS 2005 is applied to citrus seedlings grown in greenhouses or shadehouses.

One fluid ounce = 29.5 milliliters = 6 teaspoons.

Crop	Disease	Application Rate per 1 gallon water:	ppm copper per 1 gallon water	Instructions
Citrus (Non-	Brown Rot, Citrus	0.14-0.27 oz	50 -100 ppm	Begin applications when

Bearing Nursery)	Canker, Greasy Spot, Melanose, Pink Pitting, Scab	(4-8 mL) per gallon water		disease first threatens. Repeat at 30 day intervals or as needed depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	0.20-0.27 oz (6-8 mL)	75-100 ppm	Apply weekly when plants begin to vine. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	0.20 oz. (6 mL)	75 ppm	Begin applications prior to development of disease symptoms. Repeat at 7 to 10 day intervals or as needed depending on disease pressure.

Pepper	Bacterial Spot	0.20-0.27 oz (6-8 mL)	75-100 ppm	Begin applications when conditions favor disease development and repeat at 5 to 10 day intervals or as needed depending on severity. Use the higher rates when conditions favor disease.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.20-0.27 oz (6-8 mL)	75-100 ppm	Begin applications when disease first threatens and repeat at 5 to 10 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a safe place away from PETS AND KEEP OUT OF THE REACH OF CHILDREN. Store away from excessive heat. CS 2005 will freeze. Always keep container closed. Store CS 2005 in its original container only. [Bulk CS 2005 shall be stored and handled in 316L stainless steel, fiberglass, PVC's, polypropylenes or plastic equipment. Keep away from galvanized pipe and any nylon storage handling equipment.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Open dumping is prohibited. In the event of a spill, neutralize with limestone or baking soda before disposal. Concentrate may deteriorate concrete.

Container Handling: Non-refillable containers: Do not reuse or refill this container. Offer for recycling, if available. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. **For containers 5 gallons or less:** Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat two more times. **For containers of more than 5 gallons, i.e. drums:** Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand container on end and tip back and forth several times. Turn the container over on its other end and tip back and forth several times. Empty rinsate into application equipment or mix tank or store rinsate for later use or disposal. Repeat procedure two more times. **For large containers, i.e. IBC's or totes:**

Pressure washing may be an alternative. Pressure rinsing is as follows: Empty remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after flow begins to drip. Rinsing and reuse of totes is permissible.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, los alimentos ni los piensos durante el almacenamiento o la eliminación.

Almacenamiento de pesticidas: Almacenar en un lugar seguro, alejado de las mascotas y **MANTENER FUERA DEL ALCANCE DE LOS NIÑOS**. Almacenar alejado del calor excesivo. El CS 2005 se congela. Mantener siempre el envase cerrado. Almacenar el CS 2005 únicamente en su envase original. [El CS 2005 a granel debe almacenarse y manipularse en equipos de acero inoxidable 316L, fibra de vidrio, PVC, polipropileno o plástico. Mantener alejado de tuberías galvanizadas y cualquier equipo de manipulación de almacenamiento de nailon.]

Eliminación de pesticidas: Los residuos de pesticidas son extremadamente peligrosos. La eliminación inadecuada del exceso de pesticida, mezcla de pulverización o enjuague constituye una infracción de la ley federal. Si estos residuos no pueden eliminarse mediante su uso según las instrucciones de la etiqueta, comuníquese con la Agencia Estatal de Pesticidas o de Control Ambiental, o con el representante de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener orientación. Se prohíbe el vertido a cielo abierto. En caso de derrame, neutralice con piedra caliza o bicarbonato de sodio antes de desecharlo. El concentrado puede deteriorar el hormigón.

Manejo de contenedores: Envases no rellenables: No reutilice ni rellene este contenedor. Ofrézcalo para reciclaje, si está disponible. No reutilice el contenedor para contener materiales que no sean pesticidas o pesticidas diluidos (agua de enjuague). Después de vaciarlo y limpiarlo, podría permitirse almacenar temporalmente agua de enjuague u otros materiales relacionados con pesticidas en el contenedor. Comuníquese con la agencia reguladora de su estado para determinar las prácticas permitidas en su estado.

Para envases de 5 galones o menos: Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y drene 10 segundos después de que el flujo comience a gotear. Llene el envase hasta $\frac{1}{4}$ de su capacidad con agua y vuelva a taparlo. Agite durante 10 segundos. Vierta el líquido de enjuague en el equipo de aplicación o en un tanque de mezcla, o guarde el líquido de enjuague para su uso posterior o eliminación. Drene durante 10 segundos después de que el flujo comience a gotear. Repita dos veces más.

Para envases de más de 5 galones, es decir, bidones: Enjuague tres veces como se indica a continuación: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llene el envase hasta $\frac{1}{4}$ de su capacidad con agua. Cierre bien y apriete bien los cierres. Incline el envase de lado y gírelo hacia adelante y hacia atrás, asegurándose de dar al menos una vuelta completa, durante 30 segundos. Coloque el envase de pie y gírelo hacia adelante y hacia atrás varias veces. Voltee el envase sobre su otro extremo y gírelo hacia adelante y hacia atrás varias veces. Vacíe el líquido de enjuague en el equipo de aplicación o en el tanque de mezcla, o guárdelo para su uso posterior o eliminación. Repita el procedimiento dos veces más.

Para contenedores grandes, como IBC o totes, el lavado a presión puede ser una alternativa. El enjuague a presión se realiza de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y continúe drenando durante 10 segundos después de que el flujo comience a gotear. Sostenga el contenedor boca abajo sobre el equipo de aplicación o el tanque de mezcla, o recoja el líquido de enjuague para su uso posterior o eliminación. Inserte la boquilla de enjuague a presión en el lateral del contenedor y enjuague a aproximadamente 40 psi durante al menos 30 segundos. Drene durante 10 segundos después de que el flujo comience a gotear. Se permite el enjuague y la reutilización de los totes.

LIMITED WARRANTY AND LIMITATION OF REMEDIES

The directions for use of this product are believed to be reliable and must be followed carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the Seller's control. To the extent consistent with applicable laws, Buyer assumes all risks of use, handling, application, storage, or disposal of this product not in strict accordance with the use directions. To the extent consistent with applicable laws, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with the use directions. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

